

1 PATIENT INFORMATION:

Name: _____
Address: _____
City: _____ State: _____ Zip: _____
Phone: _____ Alt. Phone: _____
Email: _____
DOB: _____ Gender: M F Caregiver: _____
Height: _____ Weight: _____ Allergies: _____

2 PRESCRIBER INFORMATION:

Name: _____
Address: _____
City: _____ State: _____ Zip: _____
Phone: _____ Fax: _____
NPI: _____ DEA: _____
Tax I.D.: _____
Office Contact: _____ Phone: _____

3 STATEMENT OF MEDICAL NECESSITY: (Please Attach All Medical Documentation)

Date of Diagnosis: _____ Patient also taking Methotrexate? Yes No
ICD-10: _____ Serious or active infection present? Yes No
Other: _____ Hep B ruled out or treatment started? Yes No
TB Test: Positive Negative Date: _____ Does patient have latex allergy? Yes No
LFT: ALT: _____ AST: _____ Date: _____

Prior Failed Treatments:

Azulfidine® Celebrex® Methotrexate
 Biologics Corticosteroids Others
 Calcipotriene Indocin®

Indicate Drug Name and Length of Treatment:

If Prior Authorization is denied, recommended formulary alternatives will be provided to the prescriber based upon the patient's insurance coverage.

4 INJECTION TRAINING:

To Be Administered by Pharmacist (State of Missouri Only) Pharmacist to Provide Training Patient Trained in MD Office Manufacturer Nurse Support

5 PICK UP OR DELIVERY:

Delivery to Patient's Home Delivery to Physician's Office Pharmacy to Coordinate

6 INSURANCE INFORMATION:

Please Include Front and Back Copies of Pharmacy and Medical Card

PRESCRIPTION INFORMATION: (Please be sure to choose both induction and maintenance dose where applicable)

Patient Name: _____ Patient's Date of Birth: _____

Medication	Dosage & Strength	Direction	QTY	Refills
<input type="checkbox"/> ACTEMRA®	<input type="checkbox"/> 162mg/0.9ml Prefilled Syringe <input type="checkbox"/> 162mg/0.9ml Prefilled Autoinjector (ACTPen™)	<input type="checkbox"/> Inject 162mg SC every other week, followed by an increase to every week based on clinical response (Adults < 220 lbs) <input type="checkbox"/> Inject 162mg SC every week (Adults > 220 lbs or Patients with SJIA > 66 lbs) <input type="checkbox"/> Inject 162mg SC every 2 weeks (Patients with SJIA < 66 lbs or Patients with PJIA > 66 lbs) <input type="checkbox"/> Inject 162mg SC every 3 weeks (Patients with PJIA < 66 lbs)		
<input type="checkbox"/> CIMZIA®	<input type="checkbox"/> Prefilled Syringe Starter Kit <input type="checkbox"/> 200mg/ml Prefilled Syringe <input type="checkbox"/> 200mg Lyophilized Powder Vial	<input type="checkbox"/> Induction Dose: Inject 400mg SC on day 1, day 14 and day 28 <input type="checkbox"/> Maintenance: Inject 400mg SC every 4 weeks <input type="checkbox"/> Maintenance: Inject 200mg SC every other week	6	0
<input type="checkbox"/> COSENTYX®	<input type="checkbox"/> 150mg/ml Sensuready® Pen <input type="checkbox"/> 150mg/ml Prefilled Syringe <input type="checkbox"/> 150mg/ml Lyophilized Powder Vial	<input type="checkbox"/> Induction Dose: Inject 150mg SC at weeks 0, 1, 2, 3, and 4 <input type="checkbox"/> Induction Dose: Inject 300mg SC at weeks 0, 1, 2, 3, and 4 <input type="checkbox"/> Maintenance Dose: Inject 150mg SC every four weeks <input type="checkbox"/> Maintenance Dose: Inject 300mg SC every four weeks	5 10 1 2	0 0
<input type="checkbox"/> ENBREL®	<input type="checkbox"/> 50mg/ml Sureclick Autoinjector <input type="checkbox"/> 50mg/ml Enbrel Mini™ Prefilled Cartridge <input type="checkbox"/> For Enbrel Mini™ only: AutoTouch™ Autoinjector <input type="checkbox"/> 50mg/ml Prefilled Syringe <input type="checkbox"/> 25mg/0.5ml Prefilled Syringe <input type="checkbox"/> 25mg/ml Vial	<input type="checkbox"/> Inject 50mg SC once a week <input type="checkbox"/> Inject 25mg SC twice a week (3-4 days apart) <input type="checkbox"/> Other _____	4 1	
<input type="checkbox"/> HUMIRA®	<input type="checkbox"/> 40mg/0.4ml Pen <input type="checkbox"/> 40mg/0.4ml Prefilled Syringe <input type="checkbox"/> 40mg/0.8ml Pen <input type="checkbox"/> 40mg/0.8ml Prefilled Syringe <i>All strengths and dosages listed are Humira® Citrate Free</i>	<input type="checkbox"/> Inject 40mg SC every other week <input type="checkbox"/> Inject 40mg SC once a week <input type="checkbox"/> Patient has signed HUMIRA Complete form		
<input type="checkbox"/> KEVZARA®	<input type="checkbox"/> 150mg/1.14ml Prefilled Syringe <input type="checkbox"/> 150mg/1.14ml Prefilled Pen <input type="checkbox"/> 200mg/1.14ml Prefilled Syringe <input type="checkbox"/> 200mg/1.14ml Prefilled Pen	<input type="checkbox"/> Inject 150mg SC every 2 weeks <input type="checkbox"/> Inject 200mg SC every 2 weeks	2 2	
<input type="checkbox"/> OLUMIANT®	<input type="checkbox"/> 2mg Tablet	<input type="checkbox"/> Take 1 tablet by mouth with or without food daily	30	
<input type="checkbox"/> ORENCIA®	<input type="checkbox"/> 250mg Lyophilized Powder Vial <input type="checkbox"/> 125mg/ml ClickJect™ Autoinjector <input type="checkbox"/> 50mg/0.4ml Prefilled Syringe <input type="checkbox"/> 87.5mg/0.7ml Prefilled Syringe <input type="checkbox"/> 125mg/ml Prefilled Syringe	<input type="checkbox"/> Induction Dose: Patient Weight < 132 lbs: 500mg; 132-220 lbs: 750mg; > 220 lbs: 1000mg administered IV, then inject 125mg SC within 24 hours <input type="checkbox"/> Inject 50mg SC once a week (10 to less than 25kg) <input type="checkbox"/> Inject 87.5mg SC once a week (25 to less than 50kg) <input type="checkbox"/> Inject 125mg SC once a week (50kg or more)	4 4 4	0
<input type="checkbox"/> OTEZLA® (for PsA)	<input type="checkbox"/> Starter Pack (Titration) <input type="checkbox"/> 30mg Tablets	<input type="checkbox"/> Starter Pack: Take one tablet in the morning on day 1, then take one tablet in the morning and one tablet in the evening as directed on the starter pack <input type="checkbox"/> Maintenance: Take one 30mg tablet by mouth twice daily	1 60	0
<input type="checkbox"/> SIMPONI®	<input type="checkbox"/> 50mg/0.5ml Smartject Autoinjector <input type="checkbox"/> 50mg/0.5ml Prefilled Syringe	<input type="checkbox"/> Inject 50mg SC once a month	1	
<input type="checkbox"/> STELARA® (for PsA)	<input type="checkbox"/> 45mg/0.5ml Prefilled Syringe (for < 220 lbs) <input type="checkbox"/> 45mg/0.5ml Vial <input type="checkbox"/> 90mg/1ml Prefilled Syringe (for > 220 lbs) <input type="checkbox"/> Yes or <input type="checkbox"/> No: STELARA SELF-INJECTION: Healthcare provider certifies that patient has been trained and is eligible for self-injection	<input type="checkbox"/> Induction Dose: Inject 1 prefilled syringe SC on day 1 <input type="checkbox"/> Maintenance: Inject 1 prefilled syringe SC on day 29, and every 12 weeks thereafter	1 1	0
<input type="checkbox"/> TALTZ®	<input type="checkbox"/> 80mg/ml Single-Dose Prefilled Autoinjector <input type="checkbox"/> 80mg/ml Single-Dose Prefilled Syringe	<input type="checkbox"/> Induction Dose: Inject 160mg SC (two 80mg injections) at weeks 0 <input type="checkbox"/> Maintenance: Inject 80mg SC every 4 weeks	2 1	0
<input type="checkbox"/> XELJANZ® <input type="checkbox"/> XELJANZ® XR	<input type="checkbox"/> 5mg Tablet <input type="checkbox"/> 11mg Tablet	<input type="checkbox"/> Take one 5mg tablet by mouth twice a day <input type="checkbox"/> Take one 11mg tablet once a day	60 30	
<input type="checkbox"/> RASUVO® <input type="checkbox"/> COLCIGEL®	<input type="checkbox"/> _____ <input type="checkbox"/> _____	<input type="checkbox"/> _____ <input type="checkbox"/> _____		

PRESCRIBER SIGNATURE: I authorize pharmacy to act as my designee for initiating and coordinating insurance prior authorizations, nursing services and patient assistance programs.

Signature: _____ Date: _____ Signature: _____ Date: _____

Substitution Permitted

Dispense As Written

Prior authorization approval and insurance benefits will be determined by the payor based upon the patient's eligibility, medical necessity, and the terms of the patient's coverage, among other things. Participation in this program is not a guarantee of prior authorization or of payment.

Confidentiality Notice: This fax is intended to be delivered only to the named addressee and contains confidential information that may be protected health information under federal and state laws. If you are not the named addressee, you should not disseminate, distribute or copy this fax. Please inform the sender immediately if you have received this document in error and then destroy this document immediately.