

## BEHAVIORAL HEALTH SPECIALTY CARE PROGRAM

Phone: **833-796-6470** • Fax: **844-841-3401** 



PATIENT INFORMATION  ame:		_	BER INFORMATION:		
ddress:		Address:			
ity:	State: Zip:	City:	State: _	Zip:	
none: /			Fax:		
mail:			DEA:		
OB: Gender: O I	M OF Caregiver:	Tax I.D.:			
eight: Weight:	Allergies:	Office Contact:	Phon	ie:	
STATEMENT OF MED					
D-10:		Acute D Chronic	Prior		
ate of Diagnosis:	Contraindications: ☐ No ☐ Ye	es	Failed Treatments:	Length of	Treatment
iagnosis Procedure(s) or Laborato	ry Test(s):	_			
	Date Performed: Results:				
f Prior Authorization is denied, rec he prescriber based upon the pati		tives will be provided to			
INJECTION TRAINING					
PICK UP OR DELIVER	RY: O Delivery to Patient	e's Home O Delivery to F	Physician's Office O Pharr	nacy to Coo	rdinate
INSURANCE INFORM	ATION: Please Include F	ront and Back Copies of I	Pharmacy and Medical Car	d	
RESCRIPTION INFORM	ATION:				
tient Name:	Alloiti		MA LID LIL AUD.		formatio
LIETIL INATTIE.	Pa	itient DOB:	Must Provide All Pres	scribtion in	
		tient DOB:		-	
	Dosage & Strength		Must Provide All Pres irection	QTY	Refills
Medication	Dosage & Strength  □ 300mg Prefilled Syringe □ 400mg Prefilled Syringe	D	irection	QTY	
Medication	Dosage & Strength  ☐ 300mg Prefilled Syringe	D		QTY	
Medication  ABILIFY MAINTENA®	Dosage & Strength  □ 300mg Prefilled Syringe □ 400mg Prefilled Syringe □ 300mg Lyophilized Powder □ 400mg Lyophilized Powder	☐ Initiation: Inject 1 prefilled s	irection  Syringe IM, by a healthcare profession	QTY	
Medication  ☐ ABILIFY MAINTENA®	Dosage & Strength  □ 300mg Prefilled Syringe □ 400mg Prefilled Syringe □ 300mg Lyophilized Powder	☐ Initiation: Inject 1 prefilled s in the deltoid or gluteal mus	irection	QTY	
Medication  ABILIFY MAINTENA®  ARISTADA INITIO®	Dosage & Strength  □ 300mg Prefilled Syringe □ 400mg Prefilled Syringe □ 300mg Lyophilized Powder □ 400mg Lyophilized Powder □ 675mg Prefilled Syringe □ 441mg Prefilled Syringe	☐ Initiation: Inject 1 prefilled s in the deltoid or gluteal mus	syringe IM, by a healthcare profession cle, and take and one 30mg dose o	QTY	
Medication  ABILIFY MAINTENA®  ARISTADA INITIO®	Dosage & Strength  □ 300mg Prefilled Syringe □ 400mg Prefilled Syringe □ 300mg Lyophilized Powder □ 400mg Lyophilized Powder □ 675mg Prefilled Syringe □ 441mg Prefilled Syringe □ 662mg Prefilled Syringe □ 882mg Prefilled Syringe	Initiation: Inject 1 prefilled s in the deltoid or gluteal mus oral aripiprazole in conjunct	syringe IM, by a healthcare profession cle, and take and one 30mg dose o	QTY	
Medication  ABILIFY MAINTENA®  ARISTADA INITIO®  ARISTADA®	Dosage & Strength  □ 300mg Prefilled Syringe □ 400mg Prefilled Syringe □ 300mg Lyophilized Powder □ 400mg Lyophilized Powder □ 475mg Prefilled Syringe □ 441mg Prefilled Syringe □ 662mg Prefilled Syringe □ 882mg Prefilled Syringe □ 1064mg Prefilled Syringe	☐ Initiation: Inject 1 prefilled s in the deltoid or gluteal mus oral aripiprazole in conjunct	syringe IM, by a healthcare profession cle, and take and one 30mg dose of ion with the first Aristada injection.	QTY	
Medication  ABILIFY MAINTENA®  ARISTADA INITIO®  ARISTADA®	Dosage & Strength  □ 300mg Prefilled Syringe □ 400mg Prefilled Syringe □ 300mg Lyophilized Powder □ 400mg Lyophilized Powder □ 475mg Prefilled Syringe □ 441mg Prefilled Syringe □ 662mg Prefilled Syringe □ 882mg Prefilled Syringe □ 1064mg Prefilled Syringe □ 2mg/0.4ml Autoinjector	Initiation: Inject 1 prefilled s in the deltoid or gluteal mus oral aripiprazole in conjunct	syringe IM, by a healthcare profession cle, and take and one 30mg dose of ion with the first Aristada injection.	QTY	
Medication  ABILIFY MAINTENA®  ARISTADA INITIO®  ARISTADA®  EVZIO®	Dosage & Strength  □ 300mg Prefilled Syringe □ 400mg Prefilled Syringe □ 300mg Lyophilized Powder □ 400mg Lyophilized Powder □ 400mg Lyophilized Powder □ 675mg Prefilled Syringe □ 662mg Prefilled Syringe □ 662mg Prefilled Syringe □ 1064mg Prefilled Syringe □ 1064mg Prefilled Syringe □ 2mg/0.4ml Autoinjector □ 39mg Prefilled Syringe □ 78mg Prefilled Syringe	☐ Initiation: Inject 1 prefilled s in the deltoid or gluteal mus oral aripiprazole in conjunct	syringe IM, by a healthcare profession cle, and take and one 30mg dose of ion with the first Aristada injection.	QTY	
Medication  ABILIFY MAINTENA®  ARISTADA INITIO®  ARISTADA®  EVZIO®	Dosage & Strength  □ 300mg Prefilled Syringe □ 400mg Prefilled Syringe □ 300mg Lyophilized Powder □ 400mg Lyophilized Powder □ 400mg Lyophilized Powder □ 675mg Prefilled Syringe □ 662mg Prefilled Syringe □ 662mg Prefilled Syringe □ 1064mg Prefilled Syringe □ 1064mg Prefilled Syringe □ 2mg/0.4ml Autoinjector □ 39mg Prefilled Syringe □ 78mg Prefilled Syringe	☐ Initiation: Inject 1 prefilled s in the deltoid or gluteal mus oral aripiprazole in conjunct	syringe IM, by a healthcare profession cle, and take and one 30mg dose of ion with the first Aristada injection.	QTY	
Medication  ABILIFY MAINTENA®  ARISTADA INITIO®  ARISTADA®  EVZIO®  INVEGA SUSTENNA®	Dosage & Strength  □ 300mg Prefilled Syringe □ 400mg Prefilled Syringe □ 300mg Lyophilized Powder □ 400mg Lyophilized Powder □ 475mg Prefilled Syringe □ 441mg Prefilled Syringe □ 662mg Prefilled Syringe □ 882mg Prefilled Syringe □ 1064mg Prefilled Syringe □ 2mg/0.4ml Autoinjector □ 39mg Prefilled Syringe	Initiation: Inject 1 prefilled s in the deltoid or gluteal mustoral aripiprazole in conjunct	syringe IM, by a healthcare profession cle, and take and one 30mg dose of ion with the first Aristada injection.	QTY	
Medication  ABILIFY MAINTENA®  ARISTADA INITIO®  ARISTADA®  EVZIO®  INVEGA SUSTENNA®	Dosage & Strength  □ 300mg Prefilled Syringe □ 400mg Prefilled Syringe □ 300mg Lyophilized Powder □ 400mg Lyophilized Powder □ 400mg Lyophilized Powder □ 675mg Prefilled Syringe □ 662mg Prefilled Syringe □ 882mg Prefilled Syringe □ 1064mg Prefilled Syringe □ 2mg/0.4ml Autoinjector □ 39mg Prefilled Syringe □ 78mg Prefilled Syringe □ 117mg Prefilled Syringe □ 156mg Prefilled Syringe □ 156mg Prefilled Syringe □ 234mg Prefilled Syringe	Initiation: Inject 1 prefilled s in the deltoid or gluteal mus oral aripiprazole in conjunct	syringe IM, by a healthcare profession cle, and take and one 30mg dose of ion with the first Aristada injection.	QTY  onal f	
Medication  ABILIFY MAINTENA®  ARISTADA INITIO®  ARISTADA®  EVZIO®  INVEGA SUSTENNA®	Dosage & Strength  □ 300mg Prefilled Syringe □ 400mg Prefilled Syringe □ 300mg Lyophilized Powder □ 400mg Lyophilized Powder □ 400mg Lyophilized Powder □ 675mg Prefilled Syringe □ 662mg Prefilled Syringe □ 882mg Prefilled Syringe □ 1064mg Prefilled Syringe □ 2mg/0.4ml Autoinjector □ 39mg Prefilled Syringe □ 78mg Prefilled Syringe □ 117mg Prefilled Syringe □ 117mg Prefilled Syringe □ 156mg Prefilled Syringe □ 273mg Prefilled Syringe □ 273mg Prefilled Syringe □ 273mg Prefilled Syringe □ 410mg Prefilled Syringe □ 410mg Prefilled Syringe □ 546mg Prefilled Syringe	Initiation: Inject 1 prefilled s in the deltoid or gluteal mus oral aripiprazole in conjunct	syringe IM, by a healthcare profession cle, and take and one 30mg dose one with the first Aristada injection.	QTY  onal f	
Medication  D ABILIFY MAINTENA®  D ARISTADA INITIO®  D ARISTADA®  D EVZIO®  D INVEGA SUSTENNA®	Dosage & Strength  □ 300mg Prefilled Syringe □ 400mg Prefilled Syringe □ 300mg Lyophilized Powder □ 400mg Lyophilized Powder □ 400mg Lyophilized Powder □ 675mg Prefilled Syringe □ 662mg Prefilled Syringe □ 882mg Prefilled Syringe □ 1064mg Prefilled Syringe □ 2mg/0.4ml Autoinjector □ 39mg Prefilled Syringe □ 117mg Prefilled Syringe □ 117mg Prefilled Syringe □ 156mg Prefilled Syringe □ 156mg Prefilled Syringe □ 273mg Prefilled Syringe □ 273mg Prefilled Syringe □ 273mg Prefilled Syringe □ 410mg Prefilled Syringe □ 410mg Prefilled Syringe □ 410mg Prefilled Syringe □ 546mg Prefilled Syringe □ 546mg Prefilled Syringe	□ Initiation: Inject 1 prefilled s in the deltoid or gluteal mus oral aripiprazole in conjunct □ □ □ □ □ □ □ □ Use INVEGA TRINZA® only after patier for at least four months	syringe IM, by a healthcare professic cle, and take and one 30mg dose o ion with the first Aristada injection.	QTY  onal f	
Medication  ABILIFY MAINTENA®  ARISTADA INITIO®  ARISTADA®  EVZIO®  INVEGA SUSTENNA®  INVEGA TRINZA®	Dosage & Strength  □ 300mg Prefilled Syringe □ 400mg Prefilled Syringe □ 300mg Lyophilized Powder □ 400mg Lyophilized Powder □ 400mg Lyophilized Powder □ 675mg Prefilled Syringe □ 662mg Prefilled Syringe □ 882mg Prefilled Syringe □ 1064mg Prefilled Syringe □ 2mg/0.4ml Autoinjector □ 39mg Prefilled Syringe □ 78mg Prefilled Syringe □ 117mg Prefilled Syringe □ 156mg Prefilled Syringe □ 156mg Prefilled Syringe □ 273mg Prefilled Syringe □ 273mg Prefilled Syringe □ 410mg Prefilled Syringe □ 410mg Prefilled Syringe □ 546mg Prefilled Syringe □ 546mg Prefilled Syringe □ 549mg Prefilled Syringe □ 819mg Prefilled Syringe	□ Initiation: Inject 1 prefilled s in the deltoid or gluteal mus oral aripiprazole in conjunct □ Use INVEGA TRINZA® only after patier for at least four months □ Inject monthly, S.C., by a hear	syringe IM, by a healthcare profession cle, and take and one 30mg dose of ion with the first Aristada injection.  In this been treated with INVEGA SUSTENNA ealthcare professional in the abdomes	QTY  onal f	
Medication  ABILIFY MAINTENA®  ARISTADA INITIO®  ARISTADA®  EVZIO®  INVEGA SUSTENNA®  INVEGA TRINZA®	Dosage & Strength  □ 300mg Prefilled Syringe □ 400mg Prefilled Syringe □ 300mg Lyophilized Powder □ 400mg Lyophilized Powder □ 400mg Lyophilized Powder □ 400mg Lyophilized Powder □ 675mg Prefilled Syringe □ 662mg Prefilled Syringe □ 882mg Prefilled Syringe □ 1064mg Prefilled Syringe □ 2mg/0.4ml Autoinjector □ 39mg Prefilled Syringe □ 117mg Prefilled Syringe □ 117mg Prefilled Syringe □ 156mg Prefilled Syringe □ 156mg Prefilled Syringe □ 273mg Prefilled Syringe □ 273mg Prefilled Syringe □ 410mg Prefilled Syringe □ 546mg Prefilled Syringe □ 819mg Prefilled Syringe □ 90mg Prefilled Syringe □ 910mg Prefilled Syringe □ 120mg Prefilled Syringe	□ Initiation: Inject 1 prefilled s in the deltoid or gluteal mus oral aripiprazole in conjunct □ □ □ □ □ □ □ □ Use INVEGA TRINZA® only after patier for at least four months	syringe IM, by a healthcare profession cle, and take and one 30mg dose of ion with the first Aristada injection.  In this been treated with INVEGA SUSTENNA ealthcare professional in the abdomes	QTY  Onal  f	
Medication  □ ABILIFY MAINTENA®  □ ARISTADA INITIO®  □ ARISTADA®  □ EVZIO®  □ INVEGA SUSTENNA®  □ INVEGA TRINZA®  □ PERSERIS™	Dosage & Strength  □ 300mg Prefilled Syringe □ 400mg Prefilled Syringe □ 300mg Lyophilized Powder □ 400mg Lyophilized Powder □ 400mg Lyophilized Powder □ 400mg Lyophilized Powder □ 675mg Prefilled Syringe □ 662mg Prefilled Syringe □ 882mg Prefilled Syringe □ 1064mg Prefilled Syringe □ 2mg/0.4ml Autonipector □ 39mg Prefilled Syringe □ 17mg Prefilled Syringe □ 156mg Prefilled Syringe □ 156mg Prefilled Syringe □ 156mg Prefilled Syringe □ 273mg Prefilled Syringe □ 273mg Prefilled Syringe □ 410mg Prefilled Syringe □ 410mg Prefilled Syringe □ 410mg Prefilled Syringe □ 120mg Prefilled Syringe □ 120mg Prefilled Syringe □ 120mg Prefilled Syringe □ 125mg Vial/Kit	□ Initiation: Inject 1 prefilled s in the deltoid or gluteal must oral aripiprazole in conjunct □ □ □ □ □ □ □ □ □ □ □ □ □ □ □ □ □ □ □	syringe IM, by a healthcare profession cle, and take and one 30mg dose of ion with the first Aristada injection.  In this been treated with INVEGA SUSTENNA ealthcare professional in the abdomes	QTY  Onal  f	
ABILIFY MAINTENA®  ARISTADA INITIO®  ARISTADA®  EVZIO®  INVEGA SUSTENNA®  INVEGA TRINZA®  PERSERIS™	Dosage & Strength  □ 300mg Prefilled Syringe □ 400mg Prefilled Syringe □ 300mg Lyophilized Powder □ 400mg Lyophilized Powder □ 400mg Lyophilized Powder □ 400mg Lyophilized Powder □ 675mg Prefilled Syringe □ 662mg Prefilled Syringe □ 882mg Prefilled Syringe □ 1064mg Prefilled Syringe □ 2mg/0.4ml Autoinjector □ 39mg Prefilled Syringe □ 117mg Prefilled Syringe □ 117mg Prefilled Syringe □ 156mg Prefilled Syringe □ 156mg Prefilled Syringe □ 273mg Prefilled Syringe □ 273mg Prefilled Syringe □ 410mg Prefilled Syringe □ 546mg Prefilled Syringe □ 819mg Prefilled Syringe □ 90mg Prefilled Syringe □ 910mg Prefilled Syringe □ 120mg Prefilled Syringe	□ Initiation: Inject 1 prefilled s in the deltoid or gluteal mus oral aripiprazole in conjunct □ □ □ □ □ □ □ □ □ □ □ □ □ □ □ □ □ □ □	syringe IM, by a healthcare professic cle, and take and one 30mg dose o ion with the first Aristada injection.  In the been treated with INVEGA SUSTENNA ealthcare professional in the abdomed and powder syringes	QTY  Onal  f	
Medication  ABILIFY MAINTENA®  ARISTADA INITIO®  ARISTADA®  EVZIO®  INVEGA SUSTENNA®  INVEGA TRINZA®  PERSERIS™  RISPERDAL CONSTA®	Dosage & Strength  □ 300mg Prefilled Syringe □ 400mg Prefilled Syringe □ 300mg Lyophilized Powder □ 400mg Lyophilized Powder □ 400mg Lyophilized Powder □ 675mg Prefilled Syringe □ 662mg Prefilled Syringe □ 662mg Prefilled Syringe □ 882mg Prefilled Syringe □ 1064mg Prefilled Syringe □ 2mg/0.4ml Autonipector □ 39mg Prefilled Syringe □ 17mg Prefilled Syringe □ 117mg Prefilled Syringe □ 156mg Prefilled Syringe □ 156mg Prefilled Syringe □ 273mg Prefilled Syringe □ 273mg Prefilled Syringe □ 273mg Prefilled Syringe □ 410mg Prefilled Syringe □ 410mg Prefilled Syringe □ 410mg Prefilled Syringe □ 12.5mg Prefilled Syringe □ 120mg Prefilled Syringe □ 125mg Vial/Kit □ 37.5mg Vial/Kit □ 37.5mg Vial/Kit □ 380mg Vial/Kit	□ Initiation: Inject 1 prefilled s in the deltoid or gluteal must oral aripiprazole in conjunct □ □ □ □ □ □ □ □ □ □ □ □ □ □ □ □ □ □ □	syringe IM, by a healthcare professic cle, and take and one 30mg dose o ion with the first Aristada injection.  In the been treated with INVEGA SUSTENNA ealthcare professional in the abdomed and powder syringes	QTY  Onal  f	
Medication  ABILIFY MAINTENA®  ARISTADA INITIO®  ARISTADA®  EVZIO®  INVEGA SUSTENNA®  INVEGA TRINZA®  PERSERIS™  RISPERDAL CONSTA®	Dosage & Strength  □ 300mg Prefilled Syringe □ 400mg Prefilled Syringe □ 300mg Lyophilized Powder □ 400mg Lyophilized Powder □ 400mg Lyophilized Powder □ 675mg Prefilled Syringe □ 662mg Prefilled Syringe □ 882mg Prefilled Syringe □ 1064mg Prefilled Syringe □ 2mg/0.4ml Autoinjector □ 39mg Prefilled Syringe □ 17mg Prefilled Syringe □ 117mg Prefilled Syringe □ 156mg Prefilled Syringe □ 117mg Prefilled Syringe □ 156mg Prefilled Syringe □ 273mg Prefilled Syringe □ 273mg Prefilled Syringe □ 279mg Prefilled Syringe □ 410mg Prefilled Syringe □ 410mg Prefilled Syringe □ 120mg Prefilled Syringe □ 120mg Prefilled Syringe □ 12.5mg Vial/Kit □ 37.5mg Vial/Kit □ 380mg Vial/Kit □ 380mg Vial/Kit	Initiation: Inject 1 prefilled s in the deltoid or gluteal mus oral aripiprazole in conjunct  Use INVEGA TRINZA® only after patier for at least four months  Inject monthly, S.C., by a heafter thoroughly mixing liqui	syringe IM, by a healthcare professic cle, and take and one 30mg dose o ion with the first Aristada injection.  In the been treated with INVEGA SUSTENNA ealthcare professional in the abdomed and powder syringes	QTY  Onal of   onal onal of   onal of   onal of   onal onal of   onal onal onal of   onal onal onal onal onal onal onal onal	
Medication  □ ABILIFY MAINTENA®  □ ARISTADA INITIO®  □ ARISTADA®  □ EVZIO®  □ INVEGA SUSTENNA®  □ INVEGA TRINZA®  □ PERSERIS™  □ RISPERDAL CONSTA®  □ VIVITROL®	Dosage & Strength  □ 300mg Prefilled Syringe □ 400mg Prefilled Syringe □ 300mg Lyophilized Powder □ 400mg Lyophilized Powder □ 400mg Lyophilized Powder □ 400mg Lyophilized Powder □ 675mg Prefilled Syringe □ 662mg Prefilled Syringe □ 882mg Prefilled Syringe □ 1064mg Prefilled Syringe □ 2mg/0.4ml Autoinjector □ 39mg Prefilled Syringe □ 117mg Prefilled Syringe □ 117mg Prefilled Syringe □ 156mg Prefilled Syringe □ 156mg Prefilled Syringe □ 273mg Prefilled Syringe □ 273mg Prefilled Syringe □ 410mg Prefilled Syringe □ 410mg Prefilled Syringe □ 546mg Prefilled Syringe □ 546mg Prefilled Syringe □ 120mg Prefilled Syringe □ 120mg Prefilled Syringe □ 120mg Prefilled Syringe □ 120mg Prefilled Syringe □ 125mg Vial/Kit □ 37.5mg Vial/Kit □ 37.5mg Vial/Kit □ 380mg Vial/Kit □ 1.5mg & 3mg Mixed Blister Pa □ 3mg Capsules □ 1.5mg & 3mg Mixed Blister Pa	Initiation: Inject 1 prefilled s in the deltoid or gluteal mus oral aripiprazole in conjunct  Use INVEGA TRINZA® only after patier for at least four months  Inject monthly, S.C., by a heafter thoroughly mixing liqui	syringe IM, by a healthcare professic cle, and take and one 30mg dose o ion with the first Aristada injection.  In the been treated with INVEGA SUSTENNA ealthcare professional in the abdomed and powder syringes	QTY  Onal of   onal onal of   onal of   onal of   onal onal of   onal onal onal of   onal onal onal onal onal onal onal onal	
Medication  ABILIFY MAINTENA®  ARISTADA INITIO®  ARISTADA®  EVZIO®	Dosage & Strength  □ 300mg Prefilled Syringe □ 400mg Prefilled Syringe □ 300mg Lyophilized Powder □ 400mg Lyophilized Powder □ 400mg Lyophilized Powder □ 675mg Prefilled Syringe □ 662mg Prefilled Syringe □ 662mg Prefilled Syringe □ 882mg Prefilled Syringe □ 1064mg Prefilled Syringe □ 2mg/0.4ml Autoinjector □ 39mg Prefilled Syringe □ 17mg Prefilled Syringe □ 117mg Prefilled Syringe □ 117mg Prefilled Syringe □ 156mg Prefilled Syringe □ 156mg Prefilled Syringe □ 273mg Prefilled Syringe □ 273mg Prefilled Syringe □ 410mg Prefilled Syringe □ 410mg Prefilled Syringe □ 120mg Prefilled Syringe □ 120mg Prefilled Syringe □ 120mg Prefilled Syringe □ 125mg Vial/Kit □ 37.5mg Vial/Kit □ 37.5mg Vial/Kit □ 380mg Vial/Kit □ 1.5mg Capsules □ 1.5mg & 3mg Mixed Blister Pa □ 3mg Capsules □ 4.5mg Capsules	Initiation: Inject 1 prefilled s in the deltoid or gluteal mus oral aripiprazole in conjunct  Use INVEGA TRINZA® only after patier for at least four months  Inject monthly, S.C., by a heafter thoroughly mixing liqui	syringe IM, by a healthcare professic cle, and take and one 30mg dose o ion with the first Aristada injection.  In this been treated with INVEGA SUSTENNA ealthcare professional in the abdomed and powder syringes	QTY  Onal of   onal onal of   onal of   onal of   onal onal of   onal onal onal of   onal onal onal onal onal onal onal onal	
Medication  □ ABILIFY MAINTENA®  □ ARISTADA INITIO®  □ ARISTADA®  □ EVZIO®  □ INVEGA SUSTENNA®  □ INVEGA TRINZA®  □ PERSERIS™  □ RISPERDAL CONSTA®  □ VIVITROL®  □ VRAYLAR®	Dosage & Strength  □ 300mg Prefilled Syringe □ 400mg Prefilled Syringe □ 300mg Lyophilized Powder □ 400mg Lyophilized Powder □ 400mg Lyophilized Powder □ 400mg Lyophilized Powder □ 675mg Prefilled Syringe □ 662mg Prefilled Syringe □ 882mg Prefilled Syringe □ 1064mg Prefilled Syringe □ 1064mg Prefilled Syringe □ 1064mg Prefilled Syringe □ 17mg Prefilled Syringe □ 78mg Prefilled Syringe □ 156mg Prefilled Syringe □ 156mg Prefilled Syringe □ 156mg Prefilled Syringe □ 273mg Prefilled Syringe □ 273mg Prefilled Syringe □ 410mg Prefilled Syringe □ 546mg Prefilled Syringe □ 546mg Prefilled Syringe □ 120mg Prefilled Syringe □ 125mg Vial/Kit □ 25mg Vial/Kit □ 37.5mg Vial/Kit □ 37.5mg Vial/Kit □ 380mg Vial/Kit □ 1.5mg Capsules □ 1.5mg & 3mg Mixed Blister Pasing Capsules □ 4.5mg Capsules □ 6mg Capsules	Initiation: Inject 1 prefilled s in the deltoid or gluteal mus oral aripiprazole in conjunct  Use INVEGA TRINZA® only after patier for at least four months  Inject monthly, S.C., by a heafter thoroughly mixing liqui	syringe IM, by a healthcare professic cle, and take and one 30mg dose o ion with the first Aristada injection.  In this been treated with INVEGA SUSTENNA ealthcare professional in the abdomed and powder syringes	onal f	
Medication  □ ABILIFY MAINTENA®  □ ARISTADA INITIO®  □ ARISTADA®  □ EVZIO®  □ INVEGA SUSTENNA®  □ INVEGA TRINZA®  □ PERSERIS™ □ RISPERDAL CONSTA®  □ VIVITROL®	Dosage & Strength  □ 300mg Prefilled Syringe □ 400mg Prefilled Syringe □ 300mg Lyophilized Powder □ 400mg Lyophilized Powder □ 400mg Lyophilized Powder □ 400mg Lyophilized Powder □ 675mg Prefilled Syringe □ 662mg Prefilled Syringe □ 882mg Prefilled Syringe □ 1064mg Prefilled Syringe □ 1064mg Prefilled Syringe □ 1078mg Prefilled Syringe □ 78mg Prefilled Syringe □ 117mg Prefilled Syringe □ 117mg Prefilled Syringe □ 156mg Prefilled Syringe □ 156mg Prefilled Syringe □ 273mg Prefilled Syringe □ 273mg Prefilled Syringe □ 410mg Prefilled Syringe □ 546mg Prefilled Syringe □ 546mg Prefilled Syringe □ 120mg Prefilled Syringe □ 125mg Vial/Kit □ 25mg Vial/Kit □ 37.5mg Vial/Kit □ 37.5mg Vial/Kit □ 380mg Vial/Kit □ 1.5mg Capsules □ 1.5mg & 3mg Mixed Blister Pasing Capsules □ 4.5mg Capsules □ 4.5mg Capsules □ 6mg Capsules □ 210mg Vial □ 300mg Vial	Initiation: Inject 1 prefilled s in the deltoid or gluteal mus oral aripiprazole in conjunct  Use INVEGA TRINZA® only after patier for at least four months  Inject monthly, S.C., by a heafter thoroughly mixing liqui	syringe IM, by a healthcare professic cle, and take and one 30mg dose o ion with the first Aristada injection.  In this been treated with INVEGA SUSTENNA ealthcare professional in the abdomed and powder syringes	onal f  onal f  and the second of the second	
Medication  ABILIFY MAINTENA®  ARISTADA INITIO®  ARISTADA®  EVZIO®  INVEGA SUSTENNA®  INVEGA TRINZA®  PERSERIS™  RISPERDAL CONSTA®  VIVITROL®  VRAYLAR®  ZYPREXA® RELPREVV™	Dosage & Strength  □ 300mg Prefilled Syringe □ 400mg Prefilled Syringe □ 300mg Lyophilized Powder □ 400mg Lyophilized Powder □ 400mg Lyophilized Powder □ 400mg Lyophilized Powder □ 675mg Prefilled Syringe □ 662mg Prefilled Syringe □ 882mg Prefilled Syringe □ 1064mg Prefilled Syringe □ 1064mg Prefilled Syringe □ 1064mg Prefilled Syringe □ 17mg Prefilled Syringe □ 78mg Prefilled Syringe □ 156mg Prefilled Syringe □ 156mg Prefilled Syringe □ 156mg Prefilled Syringe □ 273mg Prefilled Syringe □ 273mg Prefilled Syringe □ 410mg Prefilled Syringe □ 546mg Prefilled Syringe □ 546mg Prefilled Syringe □ 120mg Prefilled Syringe □ 125mg Vial/Kit □ 25mg Vial/Kit □ 37.5mg Vial/Kit □ 37.5mg Vial/Kit □ 380mg Vial/Kit □ 1.5mg Capsules □ 1.5mg & 3mg Mixed Blister Pasing Capsules □ 4.5mg Capsules □ 6mg Capsules	Initiation: Inject 1 prefilled s in the deltoid or gluteal mus oral aripiprazole in conjunct  Use INVEGA TRINZA® only after patier for at least four months  Inject monthly, S.C., by a heafter thoroughly mixing liqui	syringe IM, by a healthcare professic cle, and take and one 30mg dose o ion with the first Aristada injection.  In this been treated with INVEGA SUSTENNA ealthcare professional in the abdomed and powder syringes	onal f  onal f  and the second of the second	
Medication  ABILIFY MAINTENA®  ARISTADA INITIO®  ARISTADA®  EVZIO®  INVEGA SUSTENNA®  INVEGA TRINZA®  PERSERIS™  RISPERDAL CONSTA®  VIVITROL®  VRAYLAR®  ZYPREXA® RELPREVV™	Dosage & Strength  □ 300mg Prefilled Syringe □ 400mg Prefilled Syringe □ 300mg Lyophilized Powder □ 400mg Lyophilized Powder □ 400mg Lyophilized Powder □ 400mg Lyophilized Powder □ 675mg Prefilled Syringe □ 662mg Prefilled Syringe □ 882mg Prefilled Syringe □ 1064mg Prefilled Syringe □ 1064mg Prefilled Syringe □ 1064mg Prefilled Syringe □ 17mg Prefilled Syringe □ 78mg Prefilled Syringe □ 117mg Prefilled Syringe □ 156mg Prefilled Syringe □ 156mg Prefilled Syringe □ 17mg Prefilled Syringe □ 273mg Prefilled Syringe □ 273mg Prefilled Syringe □ 410mg Prefilled Syringe □ 410mg Prefilled Syringe □ 120mg Prefilled Syringe □ 120mg Prefilled Syringe □ 120mg Prefilled Syringe □ 120mg Vial/Kit □ 37.5mg Vial/Kit □ 37.5mg Vial/Kit □ 37.5mg Vial/Kit □ 1.5mg Capsules □ 1.5mg & 3mg Mixed Blister Pasing Capsules □ 4.5mg Capsules □ 4.5mg Capsules □ 405mg Vial □ 300mg Vial □ 300mg Vial □ 300mg Vial □ 405mg Vial	Initiation: Inject 1 prefilled s in the deltoid or gluteal mus oral aripiprazole in conjunct  Use INVEGA TRINZA® only after patier for at least four months  Inject monthly, S.C., by a he after thoroughly mixing liqui	syringe IM, by a healthcare professic cle, and take and one 30mg dose of ion with the first Aristada injection.  In this been treated with INVEGA SUSTENNA ealthcare professional in the abdomed and powder syringes	QTY  Onal of   onal onal of   onal onal of   onal onal of   onal of   onal of   onal of   onal of   onal of   onal onal of   onal onal of   onal onal of   onal onal onal onal onal onal onal onal	Refills
Medication  ABILIFY MAINTENA®  ARISTADA INITIO®  ARISTADA®  EVZIO®  INVEGA SUSTENNA®  INVEGA TRINZA®  PERSERIS™  RISPERDAL CONSTA®  VIVITROL®  VRAYLAR®  ZYPREXA® RELPREVV™	Dosage & Strength  □ 300mg Prefilled Syringe □ 400mg Prefilled Syringe □ 300mg Lyophilized Powder □ 400mg Lyophilized Powder □ 400mg Lyophilized Powder □ 400mg Lyophilized Powder □ 675mg Prefilled Syringe □ 662mg Prefilled Syringe □ 882mg Prefilled Syringe □ 1064mg Prefilled Syringe □ 1064mg Prefilled Syringe □ 17mg Prefilled Syringe □ 78mg Prefilled Syringe □ 156mg Prefilled Syringe □ 156mg Prefilled Syringe □ 156mg Prefilled Syringe □ 156mg Prefilled Syringe □ 127mg Prefilled Syringe □ 127mg Prefilled Syringe □ 234mg Prefilled Syringe □ 210mg Prefilled Syringe □ 410mg Prefilled Syringe □ 410mg Prefilled Syringe □ 120mg Prefilled Syringe □ 120mg Prefilled Syringe □ 125mg Vial/Kit □ 37.5mg Vial/Kit □ 37.5mg Vial/Kit □ 380mg Vial/Kit □ 1.5mg Capsules □ 1.5mg & 3mg Mixed Blister Pa □ 3mg Capsules □ 4.5mg Vial	Initiation: Inject 1 prefilled s in the deltoid or gluteal mus oral aripiprazole in conjunct  Use INVEGA TRINZA® only after patier for at least four months  Inject monthly, S.C., by a he after thoroughly mixing liqui	syringe IM, by a healthcare professic cle, and take and one 30mg dose of ion with the first Aristada injection.  In this been treated with INVEGA SUSTENNA ealthcare professional in the abdomed and powder syringes	QTY  Onal of   onal onal of   onal onal of   onal onal of   onal of   onal of   onal of   onal of   onal of   onal onal of   onal onal of   onal onal of   onal onal onal onal onal onal onal onal	Refills